

**CYTOKINE AND
CAM ANTAGONISTS
PRIOR AUTHORIZATION FORM**
(form effective 1/8/2024)



Keystone First

PERFORMRxSM
Next Generation Pharmacy Benefits

Fax to PerformRxSM at **1-866-497-1387**, or to speak to a representative call **1-800-588-6767**.

PRIOR AUTHORIZATION REQUEST INFORMATION			
<input type="checkbox"/> New request <input type="checkbox"/> Renewal request		Total # of pages:	
Name of office contact:		Contact's phone number:	LTC facility contact/phone:
PATIENT INFORMATION			
Patient name:		Patient ID #:	DOB:
Street address:			
Apt #:	City/state/zip:		Phone:
PRESCRIBER INFORMATION			
Prescriber name:			
Specialty:		NPI:	State license #:
Street address:			
Suite #:	City/state/zip:		
Phone:	Fax:		
CLINICAL INFORMATION			
Medication requested:			
Preferred Medications:			
<input type="checkbox"/> Actemra (tocilizumab) Syringe	<input type="checkbox"/> Humira (adalimumab) 50 mg/ml Syringe	<input type="checkbox"/> Actemra (tocilizumab) Actpen	<input type="checkbox"/> Ilumya (tildrakizumab) Syringe
<input type="checkbox"/> Actemra (tocilizumab) Vial	<input type="checkbox"/> Humira(CF) (adalimumab) Pen	<input type="checkbox"/> Adalimumab-adaz(CF) 100 mg/ml Pen	<input type="checkbox"/> Inflectra (infliximab-dyyb) Vial
<input type="checkbox"/> Adalimumab-fkjp(CF) 50 mg/ml Pen	<input type="checkbox"/> Humira(CF) (adalimumab) Syringe	<input type="checkbox"/> Adalimumab-adaz(CF) 100 mg/ml Syringe	<input type="checkbox"/> Kevzara (sarilumab) Pen
<input type="checkbox"/> Adalimumab-fkjp(CF) 50 mg/ml Syringe	<input type="checkbox"/> Infliximab Vial (Janssen's unbranded infliximab)	<input type="checkbox"/> Amjevita(CF) (adalimumab-atto) 50 mg/ml Autoinjector	<input type="checkbox"/> Kevzara (sarilumab) Syringe
<input type="checkbox"/> Avsola (infliximab-axxq) Vial	<input type="checkbox"/> Kineret (anakinra) Syringe	<input type="checkbox"/> Amjevita(CF) (adalimumab-atto) 50 mg/ml Syringe	<input type="checkbox"/> Littfalo (rittecitinib) Capsule
<input type="checkbox"/> Enbrel (etanercept) Mini Cartridge	<input type="checkbox"/> Orenzia (abatacept) Clickjet	<input type="checkbox"/> Arcalyst (riloncept) Vial	<input type="checkbox"/> Olumiant (baricitinib) Tablet
<input type="checkbox"/> Enbrel (etanercept) Sureclick Pen	<input type="checkbox"/> Orenzia (abatacept) Vial	<input type="checkbox"/> Cimzia (certolizumab pegol) Syringe	<input type="checkbox"/> Orenzia (abatacept) Syringe
<input type="checkbox"/> Enbrel (etanercept) Syringe	<input type="checkbox"/> Otezla (apremilast) Tablet	<input type="checkbox"/> Cosentyx (secukinumab) Pen	<input type="checkbox"/> Remicade (infliximab) Vial
<input type="checkbox"/> Enbrel (etanercept) Vial	<input type="checkbox"/> Simponi (golimumab) Pen	<input type="checkbox"/> Cosentyx (secukinumab) Syringe	<input type="checkbox"/> Renflexis (infliximab-abda) Vial
<input type="checkbox"/> Hadlima (adalimumab-bwwd) 50 mg/ml Pushtouch	<input type="checkbox"/> Simponi (golimumab) Syringe	<input type="checkbox"/> Cyltezo(CF) (adalimumab-adbm) 50 mg/ml Pen	<input type="checkbox"/> Rinvoq ER (upadacitinib) Tablet
<input type="checkbox"/> Hadlima (adalimumab-bwwd) 50 mg/ml Syringe	<input type="checkbox"/> Taltz (ixekizumab) Autoinjector	<input type="checkbox"/> Cyltezo(CF) (adalimumab-adbm) 50 mg/ml Syringe	<input type="checkbox"/> Siliq (brodalumab) Syringe
<input type="checkbox"/> Hadlima(CF) (adalimumab-bwwd) 100 mg/ml Pushtouch	<input type="checkbox"/> Taltz (ixekizumab) Syringe	<input type="checkbox"/> Entyvio (vedolizumab) Vial	<input type="checkbox"/> Simponi Aria (golimumab) Vial
<input type="checkbox"/> Hadlima(CF) (adalimumab-bwwd) 100 mg/ml Syringe	<input type="checkbox"/> Xeljanz (tofacitinib) Tablet	<input type="checkbox"/> Hulio(CF) (adalimumab-fkjp) 50 mg/ml Pen	<input type="checkbox"/> Skyrizi (risankizumab) On-Body Injector
<input type="checkbox"/> Humira (adalimumab) 50 mg/ml Pen	<input type="checkbox"/> Xeljanz XR (tofacitinib) Tablet	<input type="checkbox"/> Hulio(CF) (adalimumab-fkjp) 50 mg/ml Syringe	<input type="checkbox"/> Skyrizi (risankizumab) Pen
	<input type="checkbox"/> Yusimry(CF) (adalimumab-aqvh) 50 mg/ml Pen	<input type="checkbox"/> Hyrimoz(CF) (adalimumab-adaz) 100 mg/ml Pen	<input type="checkbox"/> Skyrizi (risankizumab) Syringe
		<input type="checkbox"/> Hyrimoz(CF) (adalimumab-adaz) 100 mg/ml Syringe	<input type="checkbox"/> Sotykto (deucravacitinib) Tablet
		<input type="checkbox"/> Idacio(CF) (adalimumab-aacf) 50 mg/ml Pen	<input type="checkbox"/> Spevigo (spesolimab-sbzo) Vial
		<input type="checkbox"/> Idacio(CF) (adalimumab-aacf) 50 mg/ml Syringe	<input type="checkbox"/> Stelara (ustekinumab) Syringe
		<input type="checkbox"/> Ilaris (canakinumab) Vial	<input type="checkbox"/> Stelara (ustekinumab) Vial
			<input type="checkbox"/> Tremfya (guselkumab) Autoinjector
			<input type="checkbox"/> Tremfya (guselkumab) Syringe
			<input type="checkbox"/> Xeljanz (tofacitinib) Solution
			<input type="checkbox"/> Yuflyma(CF) (adalimumab-aaty) 100 mg/ml Autoinjector
			<input type="checkbox"/> Yuflyma(CF) (adalimumab-aaty) 100 mg/ml Syringe
STARTER PACK requested (strength/formulation):		MAINTENANCE product/packaging requested (strength/formulation):	
Quantity per fill:	Refills:	Quantity per fill:	Refills:
Directions:		Directions:	
Diagnosis (<i>submit documentation</i>):		Dx code (required):	Beneficiary weight:
Is the beneficiary currently being treated with the requested medication?		<input type="checkbox"/> Yes – date of last dose: _____ <i>Submit documentation.</i>	
		<input type="checkbox"/> No	
Is the requested medication prescribed by or in consultation with a specialist (e.g., rheumatologist, dermatologist, gastroenterologist, etc)?		<input type="checkbox"/> Yes <i>If prescriber is not a specialist, submit documentation of consultation.</i>	
		<input type="checkbox"/> No	

PHARMACY INFORMATION (Prescriber to identify the pharmacy that is to dispense the medication):Deliver to: Patient's Home Physician's Office Patient's Preferred Pharmacy Name:

NPI#:

Pharmacy Phone #:

Pharmacy Fax #:

 I acknowledge that the patient agrees with the pharmacy chosen for delivery of this medication.

**Complete all sections that apply to the beneficiary and this request.
Check all that apply and submit documentation for each item.**

INITIAL REQUESTS**Drug****1. Requested drug is NON-PREFERRED:**

- Tried and failed or has a contraindication or intolerance to the preferred drugs in this class approved or medically accepted for the beneficiary's condition.
List preferred medications tried: _____

2. Requested drug is OTEZLA (apremilast) or SILIQ (brodalumab):

- Was evaluated for history of prior suicide attempt, bipolar disorder, or major depressive disorder

3. Requested drug is an oral JAK inhibitor (eg, Olumiant [baricitinib], Rinvoq [upadacitinib], Xeljanz [tofacitinib]):

- Tried and failed at least one TNF blocker or another biologic as recommended in the JAK inhibitor's package labeling
 Has a contraindication or an intolerance to TNF blockers or other biologics as recommended in the JAK inhibitor's package labeling

Diagnosis**1. ALL diagnoses:**

- Screened for hepatitis B virus infection (surface antigen, surface antibody, and core antibody)
 Screened for tuberculosis

2. Adult-onset Still's disease:

- Has predominantly systemic disease:
 Has steroid-dependent disease with intent to decrease or discontinue dose of steroid with use of Cytokine and CAM Antagonist
 Tried and failed or has a contraindication or an intolerance to systemic glucocorticoids
 Has predominantly joint disease:
 Tried and failed or has a contraindication or an intolerance to conventional DMARDs (eg, MTX)

3. Alopecia areata:

- Has alopecia universalis
 Has >50% scalp involvement or alopecia totalis
 Has alopecia that causes significant disability or impaired physical, mental, or psychosocial functioning
 Has a current episode of alopecia areata that has lasted at least 6 months

4. Ankylosing spondylitis & non-radiographic axial spondyloarthritis:

- Tried and failed a 2-week trial of or has a contraindication or an intolerance to 2 different oral NSAIDs

5. Behçet's syndrome:

- Has a diagnosis of Behçet's syndrome according to current consensus guidelines
 Has recurrent oral ulcers associated with Behçet's syndrome
 Tried and failed or has a contraindication or an intolerance to a topical corticosteroid (eg, triamcinolone dental paste)
 Tried and failed a 3-month trial of or has a contraindication or an intolerance to colchicine at maximally tolerated doses

6. Crohn's disease:

- Has moderate-to-severe disease
 Has disease that is associated with high-risk or poor prognostic features
 Tried and failed to achieve remission with or has a contraindication or an intolerance to an induction course of corticosteroids
 Tried and failed to maintain remission with or has a contraindication or an intolerance to conventional immunomodulators (e.g., AZA, 6-MP, MTX)

7. Familial Mediterranean fever:

- Tried and failed a 3-month trial of or has a contraindication or an intolerance to colchicine at maximally tolerated doses

8. Gout flare:

- Tried and failed or has a contraindication or an intolerance to NSAIDs at maximally tolerated doses
 Tried and failed or has a contraindication or an intolerance to colchicine at maximally tolerated doses
 Tried and failed or has a contraindication or an intolerance to corticosteroids
 Has a medical reason why repeated courses of corticosteroids are not appropriate

9. Giant cell arteritis:

- Tried and failed or has a contraindication or an intolerance to systemic glucocorticoids
 Is at high risk for glucocorticoid-related complications
 Has steroid-dependent disease with intent to decrease or discontinue dose of steroid with use of Cytokine and CAM Antagonist

10. Hidradenitis suppurativa (HS):

- Has Hurley stage II or stage III disease
 Is a candidate for or has a history of surgical intervention for HS
 Tried and failed a 3-month trial of or has a contraindication or an intolerance to topical clindamycin
 Tried and failed or has a contraindication or an intolerance to systemic antibiotics (e.g., doxycycline, minocycline, tetracycline, clindamycin)

INITIAL REQUESTS (continued)**11. Juvenile idiopathic arthritis:**

- Has systemic disease with active systemic features
- Has disease associated with any of the following:
 - Positive anti-CCP antibodies
 - Positive rheumatoid factor
 - Presence of joint damage
 - At high risk of disabling joint damage
 - High disease activity
 - Involvement of high-risk joints (cervical spine, hip, wrist)
- Tried and failed a 3-month trial of or has a contraindication or an intolerance to conventional DMARDs (e.g., MTX)
- Has active sacroiliitis and/or enthesitis:
 - Tried and failed a 2-week trial of or has a contraindication or an intolerance to oral NSAIDs

12. Plaque psoriasis:

- Has a BSA of $\geq 3\%$ that is affected
- Has involvement of critical areas of the body (eg, skin folds, face, genitals)
- Has psoriasis that causes significant disability or impaired physical, mental, or psychosocial functioning
- Has moderate-to-severe nail disease
- Tried and failed a 4-week trial of or has a contraindication or an intolerance to topical corticosteroids
- Tried and failed an 8-week trial of or has a contraindication or an intolerance to non-steroid topical medications (e.g., anthralin, calcineurin inhibitor, tazarotene, etc)

13. Polymyalgia rheumatica:

- Tried and failed or has a contraindication or an intolerance to systemic glucocorticoids
- Has steroid-dependent disease with intent to decrease or discontinue dose of steroid with use of Cytokine and CAM Antagonist

14. Psoriatic arthritis:

- Tried and failed an 8-week trial of or has a contraindication or an intolerance to conventional DMARDs (e.g., AZA, leflunomide, MTX, SSZ)
- Has predominantly axial disease, dactylitis, and/or enthesitis
- Has severe disease
- Has comorbid moderate-to-severe nail psoriasis
- Has comorbid active inflammatory bowel disease

15. Rheumatoid arthritis:

- Tried and failed a 3-month trial of or has a contraindication or an intolerance to conventional DMARDs (e.g., AZA, leflunomide, MTX, etc.)

16. Sarcoidosis:

- Tried and failed or has a contraindication or an intolerance to systemic glucocorticoids
- Has steroid-dependent disease
- Tried and failed or has a contraindication or an intolerance to a conventional DMARD (e.g., AZA, leflunomide, MTX, mycophenolate)

17. Ulcerative colitis:

- Has moderate-to-severe disease
- Has disease associated with multiple poor prognostic factors
- Tried and failed to achieve remission with or has a contraindication or an intolerance to an induction course of corticosteroids
- Tried and failed to maintain remission with or has a contraindication or an intolerance to conventional immunomodulators (e.g., AZA, cyclosporine, 6-MP, MTX)

18. Uveitis (non-infectious):

- Has comorbid juvenile idiopathic arthritis
- Has comorbid Behçet's syndrome
- Has steroid-dependent disease with intent to decrease or discontinue dose of steroid with use of Cytokine and CAM Antagonist
- Tried and failed or has a contraindication or an intolerance to systemic, topical, intraocular, or periocular corticosteroids
- Tried and failed or has a contraindication or an intolerance to conventional systemic immunosuppressives (e.g., AZA, MTX, MMF, etc)

19. Spevigo (spesolimab) for treatment of generalized pustular psoriasis (GPP) flares:

- Has received a single dose of Spevigo (spesolimab) for current GPP flare:
 - Continues to experience moderate to severe GPP flare symptoms since the previous dose
- Has not received a dose of Spevigo (spesolimab) for current GPP flare:
 - Is experiencing a moderate to severe GPP flare that warrants rapid stabilization or improvement

20. Other diagnosis:

- List other treatments tried (including start/stop dates, dose, outcomes):

RENEWAL REQUESTS

- Experienced an improvement in disease severity or level of functioning since starting therapy with the requested medication
- Is prescribed an increased dose or more frequent administration of the requested medication that is supported by peer-reviewed medical literature or national treatment guidelines
- Requested drug is OTEZLA (apremilast) or SILIQ (brodalumab):
 - Was recently reevaluated for behavioral and mood changes

PLEASE FAX COMPLETED FORM WITH REQUIRED CLINICAL DOCUMENTATION

Prescriber signature:

Date:

Confidentiality Notice: The documents accompanying this telecopy may contain confidential information belonging to the sender. The information is intended only for the use of the individual named above. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or taking of any telecopy is strictly prohibited.